

SJSU/TP-02-23
April 2002

Do macroscopic properties dictate microscopic probabilities?

J. Finkelstein¹
Department of Physics
San José State University
San José, CA 95192, U.S.A

Abstract

Aharonov and Reznik have recently argued that the form of the probabilistic predictions of quantum theory can be seen to follow from properties of macroscopic systems. An error in their argument is identified.

¹ Participating Guest, Lawrence Berkeley National Laboratory
e-mail: JLFINKELSTEIN@lbl.gov

Aharonov and Reznik (ref. [1], hereafter AR) have recently suggested that the form of the probabilistic predictions of quantum theory can be seen to follow from properties of macroscopic systems. This suggestion is made in the context of acceptance of the non-probabilistic parts of standard quantum theory (in particular the requirement that the result of a measurement of an observable must be one of the eigenvalues of the operator corresponding to that observable), as well as the requirement that quantum predictions will indeed be probabilistic; the result that AR wish to derive is the specific form those probabilistic predictions take. The purpose of this note is not to question the standard quantum predictions which AR wish to derive from a consideration of macroscopic properties - indeed, those standard predictions are also required by Gleason's theorem [2] and of course are supported by an enormous amount of experimental evidence - but rather to examine whether these predictions can indeed be said to follow from the considerations which AR present.

AR consider a collection of N identically-prepared spin- $\frac{1}{2}$ particles, which are in the product state

$$|\Psi\rangle = |\psi\rangle_1 \otimes |\psi\rangle_2 \otimes \dots \otimes |\psi\rangle_N, \quad (1)$$

where $|\psi\rangle_r$ is the state of the r th particle which is given, for any r , by

$$|\psi\rangle_r = c_+|+\rangle_r + c_-|-\rangle_r, \quad (2)$$

and where $|+\rangle_r$ and $|-\rangle_r$ are the eigenstates of the operator $(\sigma_x)_r$. AR define M_x to be the operator for the average value of the x -component of spin, that is,

$$M_x = \left(\frac{1}{N}\right) \sum_{r=1}^N (\sigma_x)_r. \quad (3)$$

Now let f_+ denote the probability of obtaining, in a measurement of σ_x on a single particle, the outcome $+1$; the result which AR wish to prove is that f_+ is equal to $|c_+|^2$.

If N is large, then if one were to measure M_x by measuring σ_x for each particle, the law of large numbers would require that the outcome would almost certainly be close to $f_+(+1) + (1 - f_+)(-1)$. Now let $\bar{\sigma}_x$ denote $|c_+|^2(+1) + |c_-|^2(-1)$ (since $|c_-|^2 = 1 - |c_+|^2$, this means $\bar{\sigma}_x = 2|c_+|^2 - 1$). If AR can show that, in a measurement of M_x on the state $|\Psi\rangle$ the outcome would almost certainly be close to $\bar{\sigma}_x$, then they would have the result which they want.

AR begin by reminding us (see also refs. [3], [4]) that if we write

$$M_x|\Psi\rangle = \bar{\sigma}_x|\Psi\rangle + |\Delta\rangle, \quad (4)$$

then the norm of $|\Delta\rangle$ vanishes in the limit $N \rightarrow \infty$. This fact might be interpreted to mean that, for $N = \infty$, $|\Psi\rangle$ is an eigenstate of M_x with eigenvalue $\bar{\sigma}_x$, which (according to the non-probabilistic part of standard quantum mechanics) would imply that a measurement of M_x on the state $|\Psi\rangle$ would surely have the outcome $\bar{\sigma}_x$. However (as AR carefully point out), N is finite for any actual system, and so, in order to obtain the desired result for N large but finite, AR introduce a “stability assumption”; they write

One way to proceed is to make an additional assumption which seems natural for macroscopically large samples: *the results of physical experiments are stable against small perturbations.*

Having made this assumption, they then argue for their result in two ways. The first of these is based on the assertion that a small change in $|\Psi\rangle$ would make it an *exact* eigenstate of M_x . In the second way, they consider a model for the interaction which results in a measurement of M_x , and apply their assumption to the post-measurement state of the N -particle system entangled with the measurement device.

In this note I point out that the first of these ways is in fact in error, and also note that, given their “stability assumption”, it is not necessary to consider the post-measurement state at all. The first argument given by AR begins with the following passage:

... for finite large N , the operator M_x fails to be a precise eigen-operator of $|\Psi\rangle$ However, by a small modification of the state to $|\Psi\rangle + |\delta\Psi\rangle$ with magnitude $\| |\delta\Psi\rangle \| = O(1/N)$, the perturbed state does become an exact eigenstate of M_x .

However, some time ago Squires [5] showed that, as N becomes large, $|\Psi\rangle$ becomes *orthogonal* to any eigenstate of M_x . So, if we let $|M_x = m\rangle$ denote any (normalized) eigenstate of M_x , and write

$$|M_x = m\rangle = |\Psi\rangle + |\delta\Psi\rangle, \quad (5)$$

we have

$$\| |\delta\Psi\rangle \|^2 = \| |\Psi\rangle \|^2 + \| |M_x = m\rangle \|^2 - 2\text{Re}\langle M_x = m|\Psi\rangle, \quad (6)$$

and since [5] $\lim_{N \rightarrow \infty} \| \langle M_x = m|\Psi\rangle \| = 0$, this gives $\lim_{N \rightarrow \infty} \| |\delta\Psi\rangle \| = 2$, contradicting the assertion of AR that $\| |\delta\Psi\rangle \|$ is $O(1/N)$.

This result of Squires can be understood as follows: let S be a subset of the integers from 1 to N , and define

$$|S\rangle = \left[\bigotimes_{r \in S} |+\rangle_r \right] \left[\bigotimes_{r \notin S} |-\rangle_r \right] \quad (7)$$

that is, $|S\rangle$ is the product N -particle state in which the r th particle is in state $|+\rangle$ if r is included in the set S , and is in state $|-\rangle$ if r is not included in the set S . Let $n(S)$ be the number of elements of S (that is, the number of $|+\rangle$ states in eq. 7), and define

$$|k\rangle = \left(\frac{1}{N_k} \right)^{\frac{1}{2}} \sum_{n(S)=k} |S\rangle. \quad (8)$$

The number of (mutually-orthogonal) terms on the RHS of this equation is the binomial coefficient $N!/ [k!(N-k)!]$, and so $|k\rangle$ is normalized if $N_k = N!/ [k!(N-k)!]$. Note that each term in the sum in eq. 8, and hence $|k\rangle$ itself, is an eigenstate of M_x , with eigenvalue $[k(+1) + (N-k)(-1)]/N$; I will write this eigenvalue as $\lambda_k = (2k/N) - 1$.

From eqs. 1, 2, 7 and 8, it is easy to see that

$$|\Psi\rangle = \sum_{k=0}^N c_+^k c_-^{(N-k)} N_k^{\frac{1}{2}} |k\rangle, \quad (9)$$

from which it follows that

$$|\langle k|\Psi\rangle|^2 = |c_+|^{2k} |c_-|^{2(N-k)} \left(\frac{N!}{k!(N-k)!} \right). \quad (10)$$

Since $|c_-|^2 = 1 - |c_+|^2$, this shows that $|\langle k|\Psi\rangle|^2$ is a binomial distribution in k . For large values of N , the maximum of this distribution is [6] at $k = |c_+|^2 N$ (and so $\lambda_k = \bar{\sigma}_x$), and Stirling's formula then shows that this maximum value is $O(N^{-\frac{1}{2}})$. This is Squires' result; roughly speaking, this result follows from the fact that, although $|\langle k|\Psi\rangle|^2$ is indeed peaked at $k = |c_+|^2 N$, the width of this peak grows as $N^{\frac{1}{2}}$. Thus there are $N^{\frac{1}{2}}$ terms which contribute significantly to the sum in eq. 9, and so the overlap of $|\Psi\rangle$ with any one of them must fall as $N^{-\frac{1}{4}}$.

So it is not true that $|\Psi\rangle$ becomes close to an exact eigenstate of M_x as $N \rightarrow \infty$. However, it could be said to come close to an approximate eigenstate, in the following sense: suppose, for any $\epsilon > 0$, we keep only

those terms in eq. 9 in which k is within ϵN of the peak value (which is $N|c_+|^2$); that is, let $a_{\pm} = N(|c_+|^2 \pm \epsilon)$, and then define the state $|\Psi_{\epsilon}\rangle$ (up to normalization) by

$$|\Psi_{\epsilon}\rangle = \sum_{k=a_-}^{a_+} c_+^k c_-^{(N-k)} N^{\frac{1}{2}} |k\rangle. \quad (11)$$

Since each term on the RHS of eq. 11 is an eigenstate of M_x with eigenvalue λ_k , it follows from the non-probabilistic part of quantum mechanics [7] that a measurement of M_x on the state $|\Psi_{\epsilon}\rangle$ would certainly have an outcome which was in the range from λ_{a_-} to λ_{a_+} , that is, from $\bar{\sigma}_x - 2\epsilon$ to $\bar{\sigma}_x + 2\epsilon$. Also, since from eq. 10 it follows (see ref. [6]) that

$$\sum_{k=a_-}^{a_+} |\langle k|\Psi\rangle|^2 \rightarrow 1 \quad \text{as } N \rightarrow \infty; \quad (12)$$

then [8]

$$\| |\Psi\rangle - |\Psi_{\epsilon}\rangle \| \rightarrow 0 \quad \text{as } N \rightarrow \infty. \quad (13)$$

So for any $\epsilon > 0$, it is possible to define a state $|\Psi_{\epsilon}\rangle$ on which a measurement of M_x is certain to have an outcome within 2ϵ of $\bar{\sigma}_x$ and which is close to $|\Psi\rangle$ for large N . The “stability assumption” of AR does not, as stated, specify precisely how stable the results of experiments are expected to be, nor whether that stability is uniform in N . Nevertheless, if one is willing to use that assumption, one could apply it directly to the state $|\Psi\rangle$ in the same spirit as AR apply it to the post-measurement state. Since a measurement of M_x on the state $|\Psi_{\epsilon}\rangle$ is certain to yield an outcome close to $\bar{\sigma}_x$, and since $|\Psi\rangle$ is close to $|\Psi_{\epsilon}\rangle$ for large N , that assumption would seem to imply that the outcome of a measurement of M_x on the state $|\Psi\rangle$ would almost certainly be close to $\bar{\sigma}_x$.

Therefore, if one accepts the AR “stability assumption”, it is not necessary to consider the post-measurement state at all; the quantum probability rule would follow from an application of this assumption to the pre-measurement state $|\Psi\rangle$. Of course, this does not mean that one could not obtain some insight from discussing, as AR have done, the post-measurement state.

Acknowledgement: I would like to acknowledge the hospitality of the Lawrence Berkeley National Laboratory, where this work was done.

References

- [1] Y. Aharonov and B. Reznik, “How macroscopic properties dictate microscopic probabilities,” preprints quant-ph/0110093 (2001) and quant-ph/0110093v2 (2002).
- [2] A. M. Gleason, J. Math. Mech. **6**, 885 (1957).
- [3] J. B. Hartle, Amer. J. Phys. **36**, 704 (1968).
- [4] E. Farhi, J. Goldstone and S. Gutmann, Ann. of Phys. **192**, 368 (1989).
- [5] E. J. Squires, Phys. Lett. **A145**, 67 (1990).
- [6] W. Feller, *An Introduction to Probability Theory and Its Applications* (Wiley, New York, 1968), Vol. I, chapters VI and VII.
- [7] In fact, this follows from the postulate that a measurement on an eigenstate of the measured operator will surely yield the corresponding eigenvalue. Just define an operator $Q = \prod_{k=a_-}^{a_+} (M_x - \lambda_k)$; the state $|\Psi_\epsilon\rangle$ is an eigenstate of Q with eigenvalue 0, so a measurement of Q on the state $|\Psi_\epsilon\rangle$ would surely yield the value 0. But one way to measure Q is to measure M_x , so a measurement of M_x must yield one of the λ_k with k between a_- and a_+ .
- [8] Relation (13) can also be seen to follow, without invoking any properties of the binomial distribution, from the fact that the norm of $|\Delta\rangle$ (defined in eq. 4) vanishes as $N \rightarrow \infty$. And since $|\langle k|\Psi\rangle|^2$ is a binomial distribution, then $\| |\Psi\rangle - |\Psi_\epsilon\rangle \| \rightarrow 0$ faster than any inverse power of N .